

CONTENTS

	Page
Acknowledgement	ii
List of Schemes and Figures	viii
List of Tables	x
Abstract	xiii
Abstrak	xv

CHAPTER I

INTRODUCTION

1.1	General	1
1.2	Lauraceae: General appearance and morphology	2
1.3	Classification of tribes	5
1.4	The genus <i>Phoebe</i>	7
1.5	The genus <i>Dehaasia</i>	8
1.6	Forest key to common <i>Phoebe</i> and <i>Dehaasia</i> sp.	9
1.7	Medicinal value	14
1.8	Objectives of study	15

CHAPTER 2

GENERAL CHEMICAL ASPECT: ISOQUINOLINE ALKALOIDS

2.1	General Definition	16
2.2	Classification of isoquinoline alkaloids	18
2.2.1	Proaporphines	20
2.2.2	Proaporphines alkaloid from plant species	26
2.2.3	Proaporphine-tryptamine dimers	30
2.2.4	Sources of proaporphine-tryptamine dimers	37
2.2.5	Aporphines	40
2.2.6	Aporphine alkaloids and their bioassay properties	47
2.2.7	Morphinoid alkaloids	53
2.2.8	Sources of morphinoid alkaloids	57
2.2.9	Bisbenzylisoquinoline alkaloids	61
2.3	Alkaloids from genus of <i>Phoebe</i> and <i>Dehaasia</i>	65
2.4	Isoquinoline used in modern medicine	79
2.5	Conclusion	86

CHAPTER 3

RESULTS AND DISCUSSION

3.1	Introduction	87
-----	--------------	----

PART A

Isolation and structural elucidation of alkaloids from *Phoebe* species (Lauraceae)

3.2	Alkaloids from <i>Phoebe grandis</i> (Nees) Merr.	88
-----	---	----

3.2.1	Alkaloids Isolated from the leaves of <i>Phoebe grandis</i>	90
3.2.1.1	Phoebegrandine C 173	90
3.2.1.2	Phoebegrandine D 174	95
3.2.1.3	(-)-Phoebescortechiniine A 175	99
3.2.1.4	Phoebegrandine A 28	102
3.2.1.5	Phoebegrandine B 29	104
3.2.1.6	Phoebegrandine E 176	107
3.2.1.7	Tetrahydropronuciferine 177	111
3.2.1.8	Tetrahydroglaziovine 178	115
3.2.2	Alkaloids from bark of <i>Phoebe grandis</i> (Nees) Merr.	117
3.2.2.1	(-)-Grandine A 179	117
3.2.2.2	(+)-Grandine B 180	122
3.2.2.3	(+)-Grandine C 183	128
3.2.2.4	(+)-Grandine D 184	132
3.2.2.5	Boldine 97 and Norboldine 98	136
3.3	Alkaloids isolated from <i>Phoebe scortechinii</i> .	137
3.3.1	(-)-Hexahyromecambrine A 186	137
3.3.2	(-)-Norhexahyromecambrine A 187	142
3.4	Alkaloids isolated from <i>Phoebe lanceolata</i>	145
3.4.1	Alkaloids isolated from the bark of <i>Phoebe lanceolata</i>	145
3.4.1.1	Liriodenine 135	145
3.4.1.2	Boldine 97	148
3.4.1.3	Norboldine 58	151
3.4.1.4	Asimilobine 57	153
3.4.1.5	(-)-Roemerine 47	156

3.4.2	Alkaloids isolated from the leaves of <i>Phoebe lanceolata</i>	158
3.4.2.1	Laurotetanine 99	158

PART B

ISOLATION AND STRUCTURAL ELUCIDATION OF ALKALOIDS FROM *DEHAASIA* SPECIES (LAURACEAE)

3.5	Introduction	160
3.6	Alkaloids isolated from the leaves of <i>Dehaasia longipedicellata</i>	160
3.6.1	(+)-Pallidine 188	161
3.6.2	(-)-Pallidine 179	164
3.6.3	(+)-Milonine 190	166
3.6.4	(-)-8,14-Dihydrosalutaridine 191	170
3.6.5	(-)-Sinoacutine 81	171
3.7	Alkaloids isolated from the leaves of <i>Dehaasia candolleana</i>	173
3.7.1	(-)-sebiferine 71	173
3.7.2	(+)-Sebiferine 192	176
3.7.3	Perakensol 193	178
3.8	Alkaloids Isolated From <i>Dehaasia incrassata</i>	181
3.8.1	(-)-Gyrolidine 194	181
3.8.2	(-)-3', 4'-Dihydrostephasubine 198	185
3.8.3	(-)-Norstephasubine 201	189
3.8.4	Stephasubine 202	191

CHAPTER 4

4.1	Radioligand receptor binding assays	193
4.1.1	Introduction	193
4.1.2	Receptor binding methodology	193
4.1.2.1	Preparation of alkaloids	193
4.1.2.2	Radioligand and receptor binding assay for screening of alkaloids	193
4.1.2.3	Membrane preparation	195
4.1.3	Results and Discussion	196
4.2	Parasite lactate dehydrogenase test (pLDH) for anti malaria <i>in vitro</i> drug screening	202

CHAPTER 5

CONCLUSION	204
-------------------	-----

CHAPTER 6

EXPERIMENTAL

6.1	General	208
6.2	Plant material	210
6.3	Extraction of plant material	210
6.4	Separation and purification of the alkaloids	213
6.5	General spectral data	216

REFERENCES	230
-------------------	-----

LIST OF FIGURES AND SCHEMES

FIGURES

1.1	<i>Phoebe grandis</i> leaves clustered at the end of the twigs.	10
1.2	<i>Dehaasia incrassata</i> leaves and fruit with fleshy reddish stalk.	10
3.1	The main correlation of the protons in phoebegrandine C 173	92
3.2	NOESY spectrum of phoebegrandine C 173	92
3.3	¹ H NMR spectrum of phoebegrandines C 173 and D 174 in CD ₃ OD.	98
3.4	HMBC spectrum of tetrahydropronuciferine 177 .	114
3.5	¹ H NMR spectrum of grandine A 179	120
3.6	¹ H-NMR spectrum of alkaloid grandine B 180	127
3.7	Partial NOE measurements of grandine C 183 showed the configuration of C-10 and C-7a.	129
3.8	¹ H NMR spectrum of grandine C 183	130
3.9	HMQC spectrum of grandine D 184 .	135
3.10	¹ H NMR spectrum of hexahydromecambrine A 186 .	139
3.11	¹ H NMR spectrum of norhexahydromecambrine A 187 .	144
3.12	¹ H NOE differential spectrum for alkaloid 97	149
3.13	Homonuclear Multiple Bond Connectivity (HMBC) experiment of alkaloid 98 (δ H correlations).	152
3.14	¹ H spectrum for alkaloid 57 and the NOE result of irradiating OMe-1	155

3.15	The numbering of pallidine 188	162
3.16	NOE differential measurement of (+)-sebiferine 192	177
4.1	Competitive binding assay of ^3H -scopolamine in the presence of atropine ($10^{-12} - 10^{-5}$ M).	198
4.2	Competitive receptor binding of ^3H -PN200-110 in the presence of nifedipine ($10^{-11} - 10^{-6}$ M).	199

SCHEMES

1.1	Classification of Lauraceae	6
1.2	<i>Phoebe</i> and <i>Dehaasia</i> species found in Malaysia.	7
2.1	Biosynthesis of various isoquinoline alkaloids from (<i>S</i>)-norcoclaurine	19
2.2	Proaporphine alkaloids as biogenetic precursor of the aporphines	20
2.3	The sign and magnitude of the specific rotations for the proaporphines and reduced proaporphines.	22
2.4	The biogenetic pathway of aporphines.	43
2.5	The biogenetic pathway and the types of morphinan alkaloids.	55
3.1	The main fragmentation patterns of phoebegrandine E 176 .	110
3.2	Suggested fragmentation patterns of grandine A 179	121
3.3	Suggested fragmentation patterns of alkaloid B 180 .	126
3.4	Biosynthesis (+/-) sebiferine	176

LIST OF TABLES

1.1	The medicinal values of <i>Phoebe</i> species .	14
2.1	¹³ C NMR data of amuronine 1 , cryprochine 3 , isocryprochine 4 , (-)- <i>N</i> -methylecrotsparinine 5 and pronuciferine 6 in CDCl ₃ (δ in ppm)	25
2.2	Generalizations for the ¹ H-NMR chemical shifts of H-9 and 9-OMe from <i>Roemeria</i> species	32
2.3	Generalizations for the ¹ H-NMR chemical shifts of H-12 of proaporphine-tryptamine from <i>Roemeria</i> and <i>Phoebe</i> species.	33
2.4	¹³ C-NMR data of (-)-roehybridine α-N-oxide 26 and (-)-roehybridine β-N-oxide 27 in CDCl ₃ (δ in ppm)	34
2.5	UV-max of aporphines	47
2.6	Chemical constituents isolated from the species of <i>Phoebe</i> and <i>Dehaasia</i>	71
3.1	Alkaloids isolated from <i>Phoebe grandis</i> .	89
3.2	¹³ C NMR (100 MHz) and ¹ H NMR (400 MHz) data for phoebegrandine C 173	94
3.3	Comparison of ¹ H and ¹³ C NMR data of phoebegrandine C 173 and phoebegrandine D 174 .	97
3.4	¹³ C (100 MHz) and ¹ H (400 MHz) NMR Data# for alkaloid Phoebescortechiniine A 175 in CDCl ₃	101
3.5	¹³ C NMR (100 MHz) and ¹ H NMR (400 MHz) data for Phoebegrandines A 28 and B 29	106

3.6	^{13}C NMR (100 MHz) and ^1H NMR (400 MHz) data for phoebegrandine E 176	109
3.7	^{13}C (100 MHz) and ^1H (400 MHz) NMR Data [#] for tetrahydropronuciferine 177 and tetrahydroglaziovine 178 .	113
3.8	^1H (400 MHz), ^{13}C , DEPT (100 MHz) data (δ in ppm) for grandine A 179 in (CDCl_3 + two drops CD_3OD).	119
3.9	^1H NMR data of Grandine B 180 , amurolin 181 and <i>O</i> -acetylamurolin 182 in CDCl_3 (δ in ppm, <i>J</i> in Hz).	123
3.10	^1H (400 MHz), ^{13}C (100 MHz) data (δ /ppm) for grandine B 180 in CDCl_3 .	125
3.11	^1H (400 MHz), ^{13}C (100 MHz) data (δ /ppm) for grandine C 183 in CDCl_3	131
3.12	^1H (400 MHz), ^{13}C (100 MHz) data (δ /ppm) for grandine D 184 in CDCl_3	133
3.13	^1H (400 MHz) data (δ /ppm) for grandine D 184 and prooxocypochrome 6 in CDCl_3	134
3.14	^{13}C (100 MHz) and ^1H (400 MHz) NMR data for alkaloid hexahydromecambrine A 186 .	141
3.15	^{13}C (100 MHz) and ^1H (400 MHz) NMR data for alkaloid norhexahydromecambrine A 187 in CDCl_3 .	143
3.16	^{13}C NMR data of liriodenine.	147
3.17	^1H (400 MHz) and ^{13}C (100 MHz) NMR data (δ /ppm in CDCl_3) for (-)-norboldine 98 , (-)-roemerine 47 and (-)-laurotetanine 99	157
3.18	Spectral data of (+) pallidine 188 (^1H NMR, 400 MHz in CDCl_3)	163